



AMERICAN COLLEGE OF SURGEONS

Inspiring Quality:
Highest Standards, Better Outcomes



Cancer Program Practice Profile Reports (CP³R)
Bladder, Breast, Cervix, Colon, Endometrium, Gastric, Kidney, Lung, Ovary, and Rectum Cancers Diagnosed 2013 - 2016

Attention!
2016 CP3R data released on November 27, 2018

Bassett Healthcare, Cooperstown, NY

Facility Selection

Facility Measures Measures Comparison

Interpreting This Report: The estimated performance rates shown below provide your cancer program with an estimate of the proportion of patients concordant with measure criteria by diagnosis year. If appropriate the CoC Standard and benchmark compliance rate is provided. This application provides cancer programs the opportunity to examine data to determine if performance rates are representative of the care provided at the institution and to review and modify case information using the review function for the measure of interest.

Bladder Breast Cervix Colon Endometrium Gastric Kidney Lung Ovary Rectum

Save to Excel

Select Measures	Measure	CoC Std / % ^	Estimated Performance Rates (%)				Review
			2013	2014	2015	2016	
Systemic chemotherapy is administered within 4 months to day preoperatively or day of surgery to 6 months postoperatively, or it is recommended for surgically resected cases with pathologic lymph node-positive (pN1) and (pN2) NSCLC (Quality Improvement)	LCT	4.5 / 85%	100.00	50.00	100.00	100.00	
Surgery is not the first course of treatment for cN2, M0 lung cases (Quality Improvement)	LNoSurg	4.5 / 85%	87.50	100.00	100.00	85.70	
At least 10 regional lymph nodes are removed and pathologically examined for AJCC stage IA, IB, IIA, and IIB resected NSCLC (Surveillance)	10RLN	Not Applicable	53.80	25.00	41.70	27.30	

Background: The Web-based Cancer Program Practice Profile Reports (CP³R) offer local providers comparative information to assess adherence to and consideration of standard of care therapies for major cancers. This reporting tool provides a platform from which to promote continuous practice improvement to improve quality of patient care at the local level and also permits hospitals to compare their care for these patients relative to that of other providers.